

頸顏面骨壞死術後傷口照護 (92095C)

病歷記錄表（範本）

姓名:鍾xx

病歷號碼:69xxxx

日期	健保序號	治療部位	處置	醫令	醫師簽章	備註
106/03/01	000X	99	CC: (參考範例) (1) A rough object at left lower jaw for two weeks. (2) 左下齒槽尖銳物出現約2周。我有服用(或注射)雙磷酸鹽藥物/頭頸部放射治療病史	92095C	× × ×	
			PE: (參考範例) (1) A sequestrum exposure at left lower gingiva about 30 mm in size. X-ray showed osteolytic lesion at left alveolus about 30*20 mm in size. Bad odor. Tooth 35, 36, 37 missed. (2) A sharp bone exposure at left lower gingiva about 10 mm in size. Tooth 35, 36, 37 mobility grade 1. X-ray showed osteolytic lesion at left alveolus about 30*20 mm in size.			
			Dx: (參考範例) (1) osteonecrosis of left mandible. Medicine related. (2) osteonecrosis of left mandible. radiotherapy related.			
			Tx: (參考範例) (1) Partially/totally remove sequestrum under local anesthesia. Normal saline irrigation. Oral care instruction. (2) selective grinding of sharp bone under local anesthesia.. Diluted betadine solution irrigation. Oral care instruction.			

ICD-10: M87.180(MRONJ , 藥物引起之頸骨壞死)

M27.2 (Osteoradionecrosis, 放射治療相關之頸骨壞死)